

Abbreviation	Meaning	Abbreviation	Meaning
<b>A&amp;O</b>	Alert and Oriented	<b>ECG</b>	Electrocardiogram
<b>AAA</b>	Abdominal aortic aneurysm	<b>ED</b>	Emergency department
<b>ABD</b>	Abdominal	<b>EE</b>	Employee
<b>AC</b>	Antecubital	<b>EJ</b>	External jugular
<b>ACM</b>	Advanced Case Management	<b>ER</b>	Employer
<b>AED</b>	Automated external defibrillator	<b>ESRD</b>	End stage renal disease
<b>AEIOU-TIPS</b>	Alcohol, epilepsy, infection, overdose, uremia, trauma	<b>ETA</b>	Estimated Time of Arrival
<b>ALS</b>	Advanced life support	<b>ETOH</b>	Ethyl alcohol
<b>AMPLE</b>	Allergies, Medications, Prior history, Last meal eaten, Events leading up to injury/illness	<b>ETT</b>	Endotracheal tube
<b>AMS</b>	Altered mental status	<b>f/u</b>	Follow-up
<b>APRN</b>	Advanced practice registered nurse	<b>FROI</b>	First report of injury
<b>ASA</b>	Acetylsalicylic acid	<b>GCS</b>	Glasgow coma scale
<b>BG</b>	Blood glucose	<b>GI</b>	Gastrointestinal
<b>BLS</b>	Basic life support	<b>GSW</b>	Gunshot wound
<b>BM</b>	Bowel movement	<b>GU</b>	Genitourinary
<b>BP</b>	Blood pressure	<b>H&amp;T'S</b>	Hypovolemia, Hypoxia, Hydrogen ion (acidosis), Hyper-/hypokalemia, Hypoglycemia, Hypothermia – Toxins, Tamponade (cardiac), Tension pneumothorax, Thrombosis (heart/lung), Trauma
<b>BSI</b>	Body substance isolation	<b>HCTZ</b>	Hydrochlorothiazide
<b>BVM</b>	Bag valve mask	<b>HEENT</b>	Head, Ears, Eyes, nose, throat
<b>CC</b>	Chief complaint	<b>HPI</b>	History of Present Illness
<b>CCC</b>	Continuous cardiac compressions	<b>HR</b>	Heart rate
<b>CHF</b>	Congestive heart failure	<b>HSE</b>	Health & Safety Engineer
<b>CM</b>	Case management	<b>HTN</b>	Hypertension
<b>COPD</b>	Chronic obstructive pulmonary disease	<b>Hx</b>	History
<b>COS</b>	CAREonsite	<b>IA</b>	Intake agent
<b>CP</b>	Chest pain	<b>ICP</b>	Intracranial pressure
<b>CPAP</b>	Continuous positive airway pressure	<b>ILS</b>	Intermediate life support
<b>CPR</b>	Cardiopulmonary resuscitation	<b>IM</b>	Intramuscular
<b>CTS</b>	Carpal tunnel symptoms	<b>IN</b>	Intranasal
<b>CV</b>	Clinic visit	<b>IO</b>	Intraosseous
<b>CVA</b>	Cardiovascular accident	<b>IV</b>	Intravenous
<b>DC</b>	Discontinue	<b>IVP</b>	Intravenous push
<b>DCAP-BTLS</b>	Deformities, contusions, abrasions, punctures/penetrations, burns, tenderness, lacerations, swelling	<b>IVPB</b>	Intravenous piggyback
<b>DKA</b>	Diabetic ketoacidosis	<b>IW</b>	Injured worker
<b>DNR</b>	Do not Resuscitate	<b>JVD</b>	Jugular venous distention
<b>DOA</b>	Dead on arrival	<b>L / (L)</b>	Left
<b>DOB</b>	Date of birth	<b>LAC</b>	Laceration
<b>DOH</b>	Date of hire	<b>LOC</b>	Loss of Consciousness
<b>DOI</b>	Date of injury	<b>LOV</b>	Last office visit
<b>DVT</b>	Deep vein thrombosis	<b>LTCM</b>	Long term case management
<b>Dx</b>	Diagnosis	<b>LUQ</b>	Left upper quadrant
<b>MAD</b>	Mucosal atomizer device	<b>LVM</b>	Left voicemail

Abbreviation	Meaning	Abbreviation	Meaning
<b>MCO</b>	Managed care organization	<b>PVC</b>	Premature ventricular contraction
<b>MCR</b>	Mobile Care Response	<b>q</b>	Every
<b>MD</b>	Medical doctor	<b>R / (R)</b>	Right
<b>MI</b>	Myocardial infarction	<b>ROM</b>	Range of motion
<b>MOI</b>	Mechanism of Injury	<b>ROSC</b>	Return of Spontaneous Circulation
<b>MPN</b>	Medical provider network	<b>RR</b>	Respiratory rate
<b>MRSA</b>	Methicillin resistant staph aureus	<b>RTW</b>	Return to work
<b>MVA/MVC</b>	Motor vehicle accident/ Motor Vehicle Collision	<b>RUQ</b>	Right upper quadrant
<b>NC</b>	Nasal cannula	<b>Rx</b>	Prescription medication
<b>NKDA</b>	No known drug allergies	<b>S/P</b>	Status/post
<b>NOI</b>	Nature of illness	<b>S/S</b>	Signs / symptoms
<b>NOV</b>	Next office visit		Symptoms, Allergies, Medication, Prior history, Last meal eaten, Events leading up to injury/ illness
<b>NP</b>	Nurse practitioner	<b>SAMPLE</b>	
<b>NPA</b>	Nasopharyngeal airway	<b>SL</b>	Sublingual
<b>NPO</b>	Nothing by Mouth	<b>SOB</b>	Shortness of Breath
<b>NRB</b>	Non-rebreather	<b>SPo2</b>	Pulse oximetry
<b>NS</b>	Normal saline	<b>SQ</b>	Subcutaneous
<b>NSAID</b>	Non-steroidal anti-inflammatory drug	<b>SVT</b>	Supraventricular tachycardia
<b>NSR</b>	Normal sinus rhythm	<b>Sx</b>	Symptoms or surgery
<b>NTG</b>	Nitroglycerine	<b>SZ</b>	Seizure
<b>NV</b>	Nausea / vomiting	<b>TCA</b>	Tricyclic antidepressants
<b>OB</b>	Obstetrics	<b>Td</b>	Tetanus and diphtheria combined vaccine
<b>OEMSTS</b>	Office of Emergency Medical Services and Trauma Sys	<b>Tdap</b>	Tetanus, diphtheria and pertussis comb vaccine
<b>OHT</b>	Occupational Health Technician	<b>Telemed</b>	Telemedicine
<b>ON</b>	Onsite nurse	<b>TFTC</b>	Trauma field triage criteria
<b>OOW</b>	Out of work	<b>TIA</b>	Transient ischemic attack ("mini-stroke")
<b>OPA</b>	Oropharyngeal airway	<b>TKO/KVO</b>	To keep open / keep vein open
<b>OPQRST</b>	Onset, Provokes, Radiates, Severity, Time (To evaluate	<b>TM</b>	Telemedicine
<b>OSHA</b>	Occupational Safety and Health Administration	<b>TN</b>	Telephonic nurse
<b>OT</b>	Occupational therapy	<b>TNT</b>	Telephonic nurse triage
<b>OTC</b>	Over the counter	<b>TOC</b>	Transfer of Care
<b>OTL</b>	Operations Team Lead	<b>TOI</b>	Time of injury
<b>PA</b>	Physician assistant	<b>TPA</b>	Third-party administrator
<b>PCI</b>	Percutaneous coronary intervention	<b>TPD</b>	Temporary partial disability
<b>PCP</b>	Primary care provider	<b>TTD</b>	Temporary total disability
<b>PCR</b>	Patient care record / report	<b>Tx</b>	Treatment
<b>PEA</b>	Pulseless electrical activity	<b>UDT</b>	Up-to-date
<b>PERRL</b>	Pupils Equal Round Reactive to Light	<b>UR</b>	Utilization review
<b>PO</b>	By mouth	<b>VF</b>	Ventricular fibrillation
<b>PPD</b>	Permanent partial disability	<b>VM</b>	Voicemail
<b>PRICE</b>	Protect, rest, ice, compress, elevate	<b>VS</b>	Vital signs
<b>PRN</b>	As needed	<b>VT</b>	Ventricular tachycardia
<b>PT</b>	Physical therapy	<b>WC</b>	Workers compensation
<b>PTD</b>	Permanent total disability	<b>WD</b>	Weak and Dizzy
<b>PTP</b>	Primary treating physician	<b>WPW</b>	Wolff-Parkinson-White syndrome
		<b>WT</b>	Weight